



ELSEVIER

Journal of Pure and Applied Algebra 134 (1999) 211–218

---

---

JOURNAL OF  
PURE AND  
APPLIED ALGEBRA

---

---

# The product on topological Hochschild homology of the integers with mod 4 coefficients

John Rognes\*

*Department of Mathematics, University of Oslo, N-0316 Oslo, Norway*

Communicated by C.A. Weibel; received 28 October 1995

---

## Abstract

We compute the natural multiplication on mod 4 homotopy for the ring spectrum  $T(\mathbb{Z}) = THH(\mathbb{Z})$ , and its associated natural module action upon the mod 2 homotopy. © 1999 Elsevier Science B.V. All rights reserved.

*AMS Classification:* Primary: 55N45; secondary: 16E40; 19D55; 55P42

---

## 0. Introduction

Bökstedt's *topological Hochschild homology*  $THH(A) = T(A)$  of a ring  $A$  can be thought of as the spectrum defined by forming the Hochschild complex for  $A$ , but replacing  $A$  with its Eilenberg–Mac Lane spectrum  $HA$  everywhere, and replacing all tensor products with smash products of spectra. Intuitively, this gives a simplicial spectrum  $[q] \mapsto T(A)_q = HA \wedge \cdots \wedge HA$  (with  $q + 1$  factors), and although there are technical problems with such a statement, these were overcome by Bökstedt in [2]. Just like the Hochschild complex is a cyclic complex,  $T(A)$  admits a circle action, and can be given the structure of an  $S^1$ -spectrum. Its 0-simplices  $T(A)_0$  are  $HA$ , and the circle action defines an interesting map

$$\lambda: S^1_+ \wedge HA \rightarrow T(A),$$

mapping into the 1-skeleton of  $T(A)$ . When  $A$  is commutative,  $T(A)$  is a ring spectrum, so  $\pi_*T(A)$  is an algebra. The inclusion on 0-simplices  $HA \rightarrow T(A)$  is a ring map, so

---

\* Tel.: 47 22 85 58 45; Fax: 47 22 85 43 49; E-mail: rognes@math.uio.no.

$T(A)$  is a retract of  $HA \wedge T(A)$  and thus equivalent to a wedge of Eilenberg–Mac Lane spectra. We will be interested in the case when  $A$  equals the integers. Then  $\pi_*T(\mathbb{Z})$  is trivial in positive even degrees, so the algebra structure is trivial. Instead we will study the nontrivial algebra structure on its homotopy with finite coefficients.

Ultimately, we are interested in studying the trace map  $\text{tr} : K(\mathbb{Z}) \rightarrow T(\mathbb{Z})$  defined in [2] from algebraic  $K$ -theory to topological Hochschild homology, and its refinements  $\text{tr}_{p^n} : K(\mathbb{Z}) \rightarrow T(\mathbb{Z})^{C_{p^n}}$  defined in [4] for the various finite subgroups  $C_{p^n} \subset S^1$ . These refinements combine to define the *cyclotomic trace map*  $\text{trc} : K(\mathbb{Z}) \rightarrow TC(\mathbb{Z})$ , which was used in [5] to compute the (completed) algebraic  $K$ -theory of the  $p$ -adic integers for odd  $p$ . We can extend these calculations to the case  $p = 2$ , and an essential ingredient of this program is to algebraically structure the mod 2 homotopy of the fixed-point spectra  $T(\mathbb{Z})^{C_{2^n}}$  for varying  $n$ . This note provides the foundations for these calculations, by establishing the properties summarized in Theorem 3.2 below.

**1. Homotopy with finite coefficients**

Recall the *Moore spectrum*  $S^0/n$  defined by the cofiber sequence

$$S^0 \xrightarrow{n} S^0 \xrightarrow{i} S^0/n \xrightarrow{j} S^1.$$

The mod  $n$  homotopy of a spectrum  $X$  is defined as  $\pi_*(X; \mathbb{Z}/n) = \pi_*(X \wedge S^0/n)$ . Given a prime  $p$ , we let  $i_k$  and  $j_k$  denote the maps  $i$  and  $j$  above in the case  $n = p^k$ . If  $X$  is a ring spectrum and  $S^0/n$  has a product map  $S^0/n \wedge S^0/n \rightarrow S^0/n$ , then the mod  $n$  homotopy  $\pi_*(X; \mathbb{Z}/n)$  naturally inherits a product structure from the composite

$$(X \wedge S^0/n) \wedge (X \wedge S^0/n) \xrightarrow{(23)} (X \wedge X) \wedge (S^0/n \wedge S^0/n) \rightarrow X \wedge S^0/n.$$

Here (23) denotes the shuffle map. The mod  $p$  Moore spectrum admits such a product map for all odd primes  $p$ , but not for  $p = 2$ . This is because the cofiber sequence

$$S^0/2 \wedge S^0 \xrightarrow{1 \wedge 2} S^0/2 \wedge S^0 \xrightarrow{1 \wedge i_1} S^0/2 \wedge S^0/2 \xrightarrow{1 \wedge j_1} S^0/2 \wedge S^1$$

cannot be split, and so there is no (right) unital product map  $\mu_1 : S^0/2 \wedge S^0/2 \rightarrow S^0/2 \cong S^0/2 \wedge S^0$ . To see this note that  $Sq^1(a_0) = a_1$  in  $H^*(S^0/2; \mathbb{Z}/2) \cong \mathbb{Z}/2\{a_0, a_1\}$ , whence by the Cartan formula  $Sq^2(a_0 \wedge a_0) = a_1 \wedge a_1 \neq 0$  in  $H^*(S^0/2 \wedge S^0/2; \mathbb{Z}/2)$ , and so there can be no splitting. It follows easily that twice the identity map on the Moore spectrum  $S^0/2$  has the essential factorization

$$2 = i_{1\eta}j_1 : S^0/2 \xrightarrow{j_1} S^1 \xrightarrow{\eta} S^0 \xrightarrow{i_1} S^0/2,$$

up to homotopy. Hence, there is no algebra structure on  $\pi_*(S^0/2)$ , nor a natural algebra structure on  $\pi_*(X; \mathbb{Z}/2)$  for general ring spectra  $X$ . The observations go back to Barratt [1].

We are concerned with the two-primary homotopy of  $T(\mathbb{Z})$  in an equivariant sense. To be precise, we wish to study the mod 2 homotopy of the  $C_{2^n}$ -fixed points of  $T(\mathbb{Z})$

for all  $n$ , where  $C_{2^n} \subset S^1$  is the cyclic group of order  $2^n$ . As noted above  $T(\mathbb{Z})$  is a wedge of Eilenberg–Mac Lane spectra, so in fact  $1 \wedge 2 : T(\mathbb{Z}) \wedge S^0/2 \rightarrow T(\mathbb{Z}) \wedge S^0/2$  factors through  $\eta : T(\mathbb{Z}) \wedge S^1 \rightarrow T(\mathbb{Z})$ , which is inessential. Hence, the obstruction to finding a product on  $T(\mathbb{Z}) \wedge S^0/2$  vanishes in this case. But  $\eta : T(\mathbb{Z}) \wedge S^1 \rightarrow T(\mathbb{Z})$  is not equivariantly inessential, in the sense that its restriction even to the  $C_2$ -fixed sets is essential. Thus, it is not clear that  $T(\mathbb{Z}) \wedge S^0/2$  can be given a product structure in any natural equivariant way.

However,  $2\eta \simeq *$ , so four times the identity map on  $S^0/2$  is inessential. Hence, the cofiber sequence

$$S^0 \wedge S^0/2 \xrightarrow{4 \wedge 1} S^0 \wedge S^0/2 \xrightarrow{i_2 \wedge 1} S^0/4 \wedge S^0/2 \xrightarrow{j_2 \wedge 1} S^1 \wedge S^0/2$$

splits, and there is a pairing of spectra

$$m : S^0/4 \wedge S^0/2 \rightarrow S^0/2$$

inducing a (left) unital pairing of mod 4 and mod 2 homotopy. In a similar way,  $S^0/4$  admits a multiplication map

$$\mu_2 : S^0/4 \wedge S^0/4 \rightarrow S^0/4.$$

Oka proved in [9] that  $\mu_2$  can be chosen to be left and right unital and associative up to homotopy, but not homotopy commutative. The commutator satisfies

$$\mu_2 - \mu_2 \circ (12) = i_2 \circ \eta^2 \circ (j_2 \wedge j_2),$$

up to homotopy, where (12) is the twist map. Hence, if  $X$  is a ring spectrum such that  $\eta^2 : X \wedge S^2 \rightarrow X \wedge S^0$  is equivariantly inessential, then  $\pi_*(X; \mathbb{Z}/4)$  is equivariantly commutative. This will be the case with  $T(\mathbb{Z})$  since  $\eta^2$  maps to zero in each  $\pi_* T(\mathbb{Z})^{C_{2^n}}$ , so  $\pi_*(T(\mathbb{Z})^{C_{2^n}}; \mathbb{Z}/4)$  is naturally a commutative algebra for all  $n$ .

There is also a question of associativity (or transitivity) for the action  $m$ , i.e., whether the maps  $m(\mu_2 \wedge 1)$  and  $m(1 \wedge m)$  are homotopic as maps  $S^0/4 \wedge S^0/4 \wedge S^0/2 \rightarrow S^0/2$ . The associator (their difference) factors up to homotopy as

$$m(\mu_2 \wedge 1) - m(1 \wedge m) = i_1 \circ \alpha \circ (j_2 \wedge j_2 \wedge j_1)$$

for some map  $\alpha : S^3 \rightarrow S^0$ , and the obstruction to associativity is its mod 2 residue class in  $\pi_3(S^0)/2$ . For ring spectra  $X$  that are algebras over  $K(\mathbb{Z})$  this obstruction naturally vanishes, since the obstruction (which comes from the stable three-stem) is divisible by two in  $K_3(\mathbb{Z}) \cong \mathbb{Z}/48$ . Hence, in the case of  $T(\mathbb{Z})$  the action of the mod 4 homotopy of  $T(\mathbb{Z})^{C_{2^n}}$  upon the mod 2 homotopy is really associative for all  $n$ .

The homotopy Bockstein  $\delta_k = i_k j_k : S^0/2^k \rightarrow S^1 \rightarrow S^1/2^k$  acts on mod  $2^k$  homotopy. Oka also proved that  $\mu_2$  can be chosen so that  $\delta_2$  is a derivation, i.e., so that

$$\delta_2 \mu_2 = \mu_2(\delta_2 \wedge 1) + \mu_2(1 \wedge \delta_2),$$

up to homotopy. There are natural maps  $\rho : S^0/4 \rightarrow S^0/2$  (*coefficient reduction*) and  $\varepsilon : S^0/2 \rightarrow S^0/4$  (*coefficient extension*) mapping between the following horizontal cofiber sequences:

$$\begin{array}{ccccc}
 S^0 & \xrightarrow{i_2} & S^0/4 & \xrightarrow{j_2} & S^1 \\
 \parallel & & \downarrow \rho & & \downarrow \mu_2 \\
 S^0 & \xrightarrow{i_1} & S^0/2 & \xrightarrow{j_1} & S^1 \\
 \downarrow \mu_2 & & \downarrow \varepsilon & & \parallel \\
 S^0 & \xrightarrow{i_2} & S^0/4 & \xrightarrow{j_2} & S^1.
 \end{array} \tag{1.1}$$

**Lemma 1.1.** *These maps satisfy*

$$\rho\mu_2 = m(1 \wedge \rho) \quad \text{and} \quad \varepsilon m = \mu_2(1 \wedge \varepsilon),$$

*up to homotopy.*

**Proof.** Oka constructed  $m$  in [9] so as to satisfy the first formula.

The second formula also follows from [9], by the following argument. Let  $\hat{m} : S^1/2 \rightarrow S^0/4 \wedge S^0/2$  and  $\hat{\mu}_2 : S^1/4 \rightarrow S^0/4 \wedge S^0/4$  be the homotopy fiber maps of  $m$  and  $\mu_2$  respectively. Then  $\hat{\mu}_2\varepsilon = (1 \wedge \varepsilon)\hat{m}$  by Lemma 26 of [9], and there is an induced map of cofibers  $f : S^0/2 \rightarrow S^0/4$  satisfying  $f m = \mu_2(1 \wedge \varepsilon)$ . We compute

$$f = f m(i \wedge 1) = \mu_2(1 \wedge \varepsilon)(i \wedge 1) = \mu_2(i \wedge 1)(1 \wedge \varepsilon) = 1 \wedge \varepsilon$$

as maps  $S^0 \wedge S^0/2 = S^0/2 \rightarrow S^0/4 = S^0 \wedge S^0/4$ . So  $f = \varepsilon$  and the statement follows.  $\square$

Clearly,  $\mu_2$ ,  $m$ ,  $\rho$ ,  $\varepsilon$  and the  $i_k$ ,  $j_k$  and  $\delta_k$  are compatible with the corresponding pairings and operations in homology (with integral or finite coefficients), under the Hurewicz homomorphisms induced by  $S^0 \rightarrow H\mathbb{Z}$  or  $S^0/2^k \rightarrow H\mathbb{Z}/2^k$  as appropriate. In the diagram

$$\begin{array}{ccc}
 \pi_*(T(\mathbb{Z}); \mathbb{Z}/4) & \xrightarrow{h_2} & H_*(T(\mathbb{Z}); \mathbb{Z}/4) \\
 \left. \begin{array}{c} \uparrow \\ \varepsilon \\ \downarrow \end{array} \right\} \rho & & \left. \begin{array}{c} \uparrow \\ \varepsilon \\ \downarrow \end{array} \right\} \rho \\
 \pi_*(T(\mathbb{Z}); \mathbb{Z}/2) & \xrightarrow{h_1} & H_*(T(\mathbb{Z}); \mathbb{Z}/2)
 \end{array} \tag{1.2}$$

the two squares commute, and  $h_2$  and the right-hand side  $\rho$  are multiplicative. We shall see below that the horizontal maps are injective.

## 2. Mod 2 homology of $T(\mathbb{Z})$

We recall Bökstedt’s calculation of  $H_*(T(\mathbb{Z}); \mathbb{Z}/2)$  (spectrum homology) given in [3]. The skeleton filtration on  $T(\mathbb{Z})$  induces a spectral sequence

$$E_{**}^2 = HH_*(H_*(H\mathbb{Z}; \mathbb{Z}/2)) \Rightarrow H_*(T(\mathbb{Z}); \mathbb{Z}/2),$$

where  $H_*(H\mathbb{Z}; \mathbb{Z}/2) = \bar{\mathcal{A}} = \mathbb{Z}/2[\zeta_1^2, \chi\zeta_2, \chi\zeta_3, \dots]$  is contained in the dual of the Steenrod algebra  $H_*(H\mathbb{Z}/2; \mathbb{Z}/2) = \mathcal{A} = \mathbb{Z}/2[\zeta_1, \zeta_2, \zeta_3, \dots]$ . Here the  $\zeta_i$  are the Milnor generators in degrees  $2^i - 1$ , and  $\chi$  is the canonical involution on  $\mathcal{A}$  induced by the twist map (see [8]). By the Künneth theorem

$$E_{**}^2 \cong \bar{\mathcal{A}} \otimes E(1 \otimes \zeta_1^2, 1 \otimes \chi\zeta_2, 1 \otimes \chi\zeta_3, \dots),$$

where  $E(\ )$  denotes the exterior algebra on the listed generators. Letting  $[S^1]$  denote the fundamental class in  $H_1(S_+^1; \mathbb{Z}/2)$ , the circle action map  $\lambda$  induces a homomorphism

$$\sigma : \Sigma \bar{\mathcal{A}} \rightarrow H_*(T(\mathbb{Z}); \mathbb{Z}/2)$$

given by  $\sigma(x) = \lambda_*([S^1] \otimes x)$ . Then  $\sigma(x)$  maps to  $1 \otimes x$  in the spectral sequence above, and so the exterior generators above are  $\sigma(\zeta_1^2)$ ,  $\sigma(\chi\zeta_2)$ ,  $\sigma(\chi\zeta_3)$  and so on. In fact  $\sigma$  is a derivation because of the form of the second Hochschild boundary:

$$b_2(1 \otimes x \otimes y) = x\sigma(y) - \sigma(xy) + y\sigma(x).$$

As noted  $T(\mathbb{Z})$  is a  $H\mathbb{Z}$ -algebra, so the spectral sequence consists of (free)  $\bar{\mathcal{A}}$ -modules. Since  $E^2$  is multiplicatively generated by classes in filtration 1, clearly,  $d^r = 0$  for all  $r \geq 2$  and so the spectral sequence collapses at the  $E^2$ -term.

Let  $e_3 = \sigma(\zeta_1^2)$  and  $e_{2^i} = \sigma(\chi\zeta_i)$  for all  $i \geq 2$  define classes in  $H_*(T(\mathbb{Z}); \mathbb{Z}/2)$ , indexed by their degrees. Our classes differ from those of Bökstedt only in that we apply the involution  $\chi$  to eliminate the decomposable indeterminacy present in [3].

**Lemma 2.1.** *In  $H_*(T(\mathbb{Z}); \mathbb{Z}/2)$  (i)  $e_3^2 = 0$ , (ii)  $e_{2^i}^2 = e_{2^{i+1}}$  for all  $i \geq 2$ , (iii)  $\beta_1(e_4) = e_3$ , and (iv)  $\hat{Q}^4(e_3) = 0$ .*

**Proof.**  $T(\mathbb{Z})$  is an  $E_\infty$  ring spectrum, so its homology  $H_*(T(\mathbb{Z}); \mathbb{Z}/2)$  admits multiplicative Dyer–Lashof operations  $\hat{Q}^r$  (see [6] or [7]). Likewise, there are additive Dyer–Lashof operations  $Q^r$  on  $H_*(H\mathbb{Z}; \mathbb{Z}/2) = \bar{\mathcal{A}}$ , which are recorded in [10, Ch. III, Theorem 2.2]. In particular,  $Q^{2^i}(\chi\zeta_i) = \chi\zeta_{i+1}$ . By Bökstedt’s lemma [3, Lemma 2.9] these operations are compatible under  $\sigma$ , i.e.,  $\hat{Q}^r(\sigma(x)) = \sigma(Q^r(x))$  for all  $x$ . Hence, we can compute

$$e_3^2 = \hat{Q}^3(e_3) = \sigma(Q^3(\zeta_1^2)) = 0$$

by the Cartan formula. Similarly for  $i \geq 2$

$$e_{2^i}^2 = \hat{Q}^{2^i}(e_{2^i}) = \sigma(Q^{2^i}(\chi\zeta_i)) = \sigma(\chi\zeta_{i+1}) = e_{2^{i+1}}$$

as claimed. Next,  $\chi\zeta_2 = \zeta_2 + \zeta_1^3$ ,  $\beta_1(\zeta_1) = 1$  and  $\beta_1(\zeta_i) = 0$  for  $i \geq 2$ . Hence,

$$\beta_1(e_4) = \beta_1(\sigma(\chi\zeta_2)) = \sigma(\beta_1\zeta_2 + \beta_1\zeta_1^3) = \sigma(\zeta_1^2) = e_3.$$

Finally,

$$\hat{Q}^4(e_3) = \sigma(Q^4(\zeta_1^2)) = \sigma(Q^2(\zeta_1)^2) = 0,$$

using the Cartan formula and that  $\sigma$  is a derivation.  $\square$

Thus, as algebras and free  $\mathcal{A}$ -modules we have

$$H_*(T(\mathbb{Z}); \mathbb{Z}/2) \cong \mathcal{A}[e_3, e_4]/(e_3^2 = 0).$$

Since  $T(\mathbb{Z})$  is a wedge of Eilenberg–Mac Lane spectra the integral and mod  $2^k$  Hurewicz homomorphisms

$$h : \pi_*T(\mathbb{Z}) \rightarrow H_*T(\mathbb{Z}) \quad \text{and} \quad h_k : \pi_*(T(\mathbb{Z}); \mathbb{Z}/2^k) \rightarrow H_*(T(\mathbb{Z}); \mathbb{Z}/2^k)$$

are all injective. In the mod 2 case the image consists of  $\mathcal{A}$ -module generators. There is at most one of these in each degree, so under  $h_1$  we can identify

$$\pi_*(T(\mathbb{Z}); \mathbb{Z}/2) \cong \mathbb{Z}/2[e_3, e_4]/(e_3^2 = 0).$$

The algebra structure here is that inherited by  $\pi_*(T(\mathbb{Z}); \mathbb{Z}/2)$  as a subgroup of  $H_*(T(\mathbb{Z}); \mathbb{Z}/2)$  which turns out to be closed under multiplication, and thus becomes a subalgebra. This algebraically defined product does not need a priori to be well related to a spectrum level product on  $T(\mathbb{Z}) \wedge S^0/2$ . We shall prove instead that it is well related to the natural pairings on homotopy induced by  $\mu_2$  and  $m$ .

**Remark 2.2.** From Lemma 2.1(iv) the higher Bockstein structure on  $\pi_*(T(\mathbb{Z}); \mathbb{Z}/2)$  may be deduced. Combining these results with the corresponding odd primary calculations, Bökstedt obtained the integral answer

$$\pi_*T(\mathbb{Z}) = \begin{cases} \mathbb{Z} & \text{for } * = 0, \\ \mathbb{Z}/n & \text{for } * = 2n - 1, \\ 0 & \text{otherwise.} \end{cases}$$

### 3. Homotopy of $T(\mathbb{Z})$ with finite coefficients

We recall the additive structure on  $\pi_*(T(\mathbb{Z}); \mathbb{Z}/4)$ . The mod 2 homotopy Bockstein  $\delta_1$  is a derivation, because it is detected in homology where it corresponds to the mod 2 homology Bockstein ( $\beta_1$  dual to  $Sq^1$ ), which is a derivation. From  $\beta_1(e_4) = e_3$  we have  $\pi_3T(\mathbb{Z})_2^\wedge = \mathbb{Z}/2$ . In general,  $\pi_{4k-1}T(\mathbb{Z})_2^\wedge$  is  $\mathbb{Z}/2$  for  $k$  odd, and cyclic of order 4 or

more for  $k$  even. So

$$\pi_*(T(\mathbb{Z}); \mathbb{Z}/4) \cong \begin{cases} \mathbb{Z}/4 & \text{for } * \equiv 0, 7 \pmod{8}, \\ \mathbb{Z}/2 & \text{for } * \equiv 3, 4 \pmod{8}, \\ 0 & \text{otherwise.} \end{cases}$$

We let  $f_i \in \pi_i(T(\mathbb{Z}); \mathbb{Z}/4)$  denote an additive generator for each  $i \equiv 0, 3 \pmod{4}$ . There is a choice of sign in  $f_i$  for  $i \equiv 0, 7 \pmod{8}$ , which we will specify shortly. The classes  $f_i$  in odd degrees are integral. We note that  $2f_3 = 2f_4 = 0$ .

The homomorphisms  $\rho$  and  $\varepsilon$  linking  $\pi_*(T(\mathbb{Z}); \mathbb{Z}/2)$  and  $\pi_*(T(\mathbb{Z}); \mathbb{Z}/4)$  are easily determined using the maps of cofiber sequences (1.1):

$$\rho(f_{8k+i}) = \begin{cases} e_4^{2k} & \text{for } i = 0, \\ e_3 e_4^{2k} & \text{for } i = 3, \\ 0 & \text{for } i = 4, \\ e_3 e_4^{2k+1} & \text{for } i = 7 \end{cases}$$

and

$$\varepsilon(e_3^a e_4^{2k+b}) = \begin{cases} 2f_{8k} & \text{for } a = 0, b = 0, \\ 0 & \text{for } a = 1, b = 0, \\ f_{8k+4} & \text{for } a = 0, b = 1, \\ 2f_{8k+7} & \text{for } a = 1, b = 1. \end{cases}$$

The mod 2 Bockstein satisfies  $\delta_2(f_4) = f_3$ ,  $\delta_2(f_8) = \pm f_7$  and is a derivation. We agree to choose  $f_7$  so that  $\delta_2(f_8) = f_7$ .

We also let  $g_i \in \pi_i T(\mathbb{Z})$  denote an integral generator for each  $i \equiv 3 \pmod{4}$ , reducing mod 4 to  $f_i$  and mod 2 to  $e_i$ .

**Lemma 3.1.** *The product on the mod 4 homotopy of  $T(\mathbb{Z})$  satisfies the following relations:*

$$\begin{aligned} f_3 f_3 &= 0, & f_3 f_4 &= 2f_7, & f_3 f_7 &= 0, & f_3 f_8^k &= f_{8k+3}, \\ f_4 f_4 &= 0, & f_4 f_7 &= 0, & f_4 f_8^k &= f_{8k+4}, \\ f_7 f_7 &= 0, & f_7 f_8^k &= \pm f_{8k+7} & \text{and} & f_8^k &= \pm f_{8k} \end{aligned}$$

for all  $k \geq 0$ .

**Proof.** The claims  $f_3^2 = 0$ ,  $f_3 f_7 = 0$  and  $f_7^2 = 0$  are clear since the products lie in trivial groups.

For  $f_3 f_4$  note that  $f_3$  is integral and  $\varepsilon$  is linear with respect to multiplication by integral classes. So  $f_3 f_4 = g_3 f_4 = g_3 \varepsilon(e_4) = \varepsilon(g_3 e_4) = 2f_7$ . Likewise,  $f_4 f_7 = f_4 g_7 = \varepsilon(e_4) g_7 = \varepsilon(e_4 g_7) = \varepsilon(e_3 e_4^2) = 0$ .

Because  $\delta_2$  is a derivation  $\delta_2(f_4^2) = 2f_3 f_4 = 0$ , and since  $\delta_2$  is injective in degree eight it follows that  $f_4^2 = 0$ .

Next  $f_3 f_8^k$  is nonzero because its image under  $\rho h_2$  (which is multiplicative) equals  $e_3 e_4^{2k}$  which is nonzero. Compare with diagram (1.2). So  $f_3 f_8^k$  is the only nonzero class

in degree  $8k + 3$ , which is the generator  $f_{8k+3}$ . Likewise,  $f_7 f_8^k$  is nonzero because its image under  $\rho h_2$  equals  $e_3 e_4^{2k+1}$ , and  $f_8^k$  is nonzero because its image  $e_4^{2k}$  is also nonzero.

Finally, we conclude that  $f_4 f_8^k$  is nonzero by computing  $\delta_2(f_4 f_8^k) = f_3 f_8^k + k f_4 f_7 f_8^{k-1} = f_3 f_8^k$  which is nonzero.  $\square$

Hence, we may take  $f_i f_8^k$  as the generator in degree  $8k + i$  for all  $k \geq 0$  and  $i = 0, 3, 4, 7$ . Modulo the sign of  $f_8$ , this precisely pins down a set of generators for  $\pi_*(T(\mathbb{Z}); \mathbb{Z}/4)$ . The sign of  $f_8$  can be fixed by insisting that  $h_2(f_8) = P(e_4)$  where  $P$  is the Pontrjagin squaring operation (see [6] or [7]). We omit the details of this argument.

**Theorem 3.2.** (i) *The mod 4 homotopy algebra of  $T(\mathbb{Z})$  is*

$$\pi_*(T(\mathbb{Z}); \mathbb{Z}/4) \cong \mathbb{Z}/4[f_3, f_4, f_7, f_8] / \sim$$

with the relations  $2f_3 = 2f_4 = 0$ ,  $f_3 f_4 = 2f_7$ , and  $f_i f_j = 0$  for all other  $i, j < 8$ .

(ii) *The pairing*

$$\pi_*(T(\mathbb{Z}); \mathbb{Z}/4) \otimes \pi_*(T(\mathbb{Z}); \mathbb{Z}/2) \rightarrow \pi_*(T(\mathbb{Z}); \mathbb{Z}/2)$$

induced by  $m$  takes  $x \otimes y$  to  $\rho(x)y$  in the subalgebra product on  $\pi_*(T(\mathbb{Z}); \mathbb{Z}/2)$  inherited from  $H_*(T(\mathbb{Z}); \mathbb{Z}/2)$ . Thus,  $m(x \otimes y) = \rho(x) \cdot y$ .

(iii) *The coefficient reduction  $\rho: \pi_*(T(\mathbb{Z}); \mathbb{Z}/4) \rightarrow \pi_*(T(\mathbb{Z}); \mathbb{Z}/2)$  is an algebra homomorphism when mapping to the subalgebra product. The coefficient extension  $\varepsilon: \pi_*(T(\mathbb{Z}); \mathbb{Z}/2) \rightarrow \pi_*(T(\mathbb{Z}); \mathbb{Z}/4)$  is a  $\pi_*(T(\mathbb{Z}); \mathbb{Z}/4)$ -module homomorphism with respect to the actions induced by  $m$  and  $\mu_2$ .*

**Proof.** Part (i) summarizes Lemma 3.1 and the conventions following it.

To prove  $m(x \otimes y) = \rho(x)y$  in  $\pi_*(T(\mathbb{Z}); \mathbb{Z}/2)$  it suffices to apply the injection  $h_1$  and note that  $h_1 m(x \otimes y) = \rho h_2(x) h_1(y) = h_1(\rho(x)y)$ .

The remaining claims follow from Lemma 1.1.  $\square$

## References

- [1] M.G. Barratt, Track groups (II), Proc. London Math. Soc. (3) 5 (1951) 285–329.
- [2] M. Bökstedt, Topological Hochschild homology, Topology, to appear.
- [3] M. Bökstedt, The topological Hochschild homology of  $\mathbb{Z}$  and  $\mathbb{Z}/p$ , Ann. of Math, to appear.
- [4] M. Bökstedt, W.C. Hsiang, I. Madsen, The cyclotomic trace and algebraic  $K$ -theory of spaces, Invent. Math. 11 (1993) 465–540.
- [5] M. Bökstedt, I. Madsen, Topological cyclic homology of the integers, Asterisque 226 (1994) 57–143.
- [6] I. Madsen, Higher torsion in  $SG$  and  $BSG$ , Math. Z. 143 (1975) 55–80.
- [7] J.P. May, The homology of  $E_\infty$  spaces, The Homology of Iterated Loop Spaces, Lecture Notes in Mathematics, vol. 533, Springer, Berlin, 1976, pp. 1–68.
- [8] J. Milnor, The Steenrod algebra and its dual, Ann. of Math. 67 (1958) 150–171.
- [9] S. Oka, Multiplications on the Moore spectrum, Mem. Fac. Sci. Kyushu Univ. Ser. A 38 (1984) 257–276.
- [10] M. Steinberger, Homology operations for  $H_\infty$  and  $H_n$  ring spectra,  $H_\infty$  ring spectra and their applications, Lecture Notes in Mathematics, vol. 1176, Springer, Berlin, 1986, pp. 56–87.